



The Fatigue Paradox: A Dual-Perspective Analysis of In-Season Neuromuscular Integrity and Load Management in the National Basketball Association

Brian Serrano*
DC, PhD, ATC, CSCS, CPSS, RSCC

Chicago Bulls Basketball Club

Received Date: February 09, 2025; **Accepted Date:** February 19, 2026; **Published Date:** February 26, 2026;

***Corresponding author:** Brian Serrano, DC, PhD, ATC, CSCS, CPSS, RSC, ¹Chicago Bulls Basketball Club;
Email: brianserrano171@gmail.com

Abstract

The National Basketball Association (NBA) competitive season presents a unique physiological environment characterized by a dense game schedule, frequent travel between time zones, and a high volume of external stress via game mechanical workloads. This paper explores the Fatigue Paradox—the conflict between the necessity of proper recovery and the deleterious risk of in-season deconditioning. By synthesizing recent longitudinal data, the author examines the "devil's advocate" position that rest is the primary variable for injury prevention without the need for in-season strength and conditioning. The author then provides a counterargument centered on themes such as "Tissue Robustness" and "Braking Performance".

Introduction: The Load Profile of Modern Professional Basketball

The modern NBA pace is the fastest it has ever been, this evolution has fundamentally altered the physiological profile of the game (Huyghe et al., 2022). Players now experience a higher frequency of high-stress transitions compared to the isolation-heavy era of the early 2000s. An NBA guard can expect to perform over

100 high-magnitude accelerations ($>3.0\text{m/s}^2$) and an even higher volume of decelerations ($<-3.0\text{m/s}^2$) per game (Mandic et al., 2019). These altered physiological demands have changed the stress profiles on tissues and joints.

The primary challenge for the performance staff is the 82-game season in 25 weeks. This is further made difficult since unlike other sports which may play at set frequencies, Basketball plays an average of 3.4 games every week. In this mix are back-to-back games, 4 games in 5 night, and teams rarely have more than two full days before games. Traditional periodization models may not be optimal in this environment because the competition phase lasts six months without a true unloading period (Bartolomei et al., 2014). Looking at the origins of periodization theories, they were built on Olympic quadrennials (4-year cycles) which may not fit this context. Consequently, the performance ceiling often drops as the season progresses from a muscular strength and tissue quality standpoint, leading to a state where the athlete's mechanical capacity is exceeded by game demands—one primary driver of non-contact soft tissue injuries.

The Case Against In-Season Loading: The Passive Recovery Theory

The argument for Load Management (in any facet) is built on the premise that fatigue is cumulative and non-linear (Chiu & Barnes, 2003). These two ideas are fairly novel within the load management literature. Fatigue is cumulative, that is magnified when breaks are few and far between. For example, performance staffs often approach the schedule in blocks, separated by two days on non-games. A game block can be comprised on 4-8 games, which highlights how each game can have a compounding effect on overall fatigue. Also, moving from the traditional view of acute: chronic workload ratio which takes the average of 7 days to the average of 28 days to give a ratio, more recent evidence points to an alternate exponentially weighted model (EWMA) which uses lambda to place more emphasis on recent work (previous 7 days) over the more chronic loads (28 days) (Williams et al., 2017).

- 1. The IQVIA/NBA Longitudinal Findings (2024)** In early 2024, the NBA released findings from a nine-season study involving 150 players. The data showed no statistically significant correlation between games missed for rest and a decrease in the likelihood of future injury. This suggests that fatigue is not just a result of increased game minutes, but a failure of the athlete to adapt to those minutes. Additionally, with technologies like Second Spectrum and Kinexon to monitor external game loads, performance staffs can begin building athletic profiles. This shows that while the total game load is important, how the player got there in each individual metric is just as important. This is where performance staffs in the NBA have been working to change the traditional view of recovery between games. Recovery is not rest in the traditional sense which stresses passive modalities. For example, while recovery can be a day of complete rest, this should not be the absolute within the NBA season because this is where de-training can occur. As will be later explored in this paper, there needs to be proper undulation of court work (games, training) with strength and conditioning work to prevent detraining.
- 2. The Interference Effect and the CNS** Heavy resistance training (>85% 1RM) relies primarily on Type 2 motor unit recruitment. During a dense schedule, the Central Nervous System (CNS) often experiences neural thinning, where the speed of the signal from the motor cortex to the muscle slows

down. Put another way, the physical demands and increased cognitive load place may lead to CNS fatigue (Moreira et al., 2018). Adding heavy lifting during this phase has the potential to exacerbate this CNS fatigue. This can be perceived as part of the interference effect, where the body is unable to simultaneously recover from game-day metabolic stress and adapt to the weight room's mechanical stress.

Methodology: Quantifying the Invisible Load

To manage this paradox, the author utilized a data-centric methodology that moves beyond minutes played and into total mechanical load, akin to common external load metrics.

1. Wearable Technology and LPS Integration Using Kinexon or Catapult LPS systems, we track **External Load** via:

- **Deceleration as a Proxy:** The total sum of decelerations since research shows decelerations are stressful movements on the body. While accelerations may give us a physical load estimate, decelerations may give us a physiological load estimate.
- **Mechanical Load:** A derivative of accelerometer data that quantifies the cost of movement which is simple to understand since it uses time as the primary collecting variable.

The Paradox of The Strength and Conditioning Program

Since both longitudinal and cross-sectional data suggests complete rest doesn't prevent injury nor help increase recovery as a standalone intervention, the focus should shift to robustness and resiliency.

- 1. Tissue Stress During Games:** Deceleration is an eccentric-dominant activity, and one that is performed many times during a game. Eccentric muscle actions are more mechanically efficient but cause more structural tissue damage due to assistance from passive structures (Vasquez-Guerrero et al., 2018). As the season progresses, players often lose objective metrics such as Eccentric Impulse, meaning they perform less work in the same/more amount of time. This can be seen either in external load metrics using deceleration zones or in CMJ testing with temporal based metrics (Bishop et al., 2023). This increased braking time

increases the time joints are under tension and may lead to compensatory movement patterns, all of which may increase the risk for tendon and soft tissue injury.

2. **Micro-Dosing as a Biological Reset:** Rather than traditional 60-minute training sessions, this method utilizes much shorter 15-20 minute micro-doses focused on providing a necessary stimulus during the season. However, the focus is on a minimum effective dose without extraneous movements or exercises. This may look different based on athletes needs but typically involves a compound movement for the upper and lower body, core, and 1-2 accessory movements.

Practical Application: Why Tissue Capacity Matters

This is where the performance staff (led by the strength and conditioning coaches) can use education and implementation to show athletes and at times the organization the value of training during the season. For example, we understand that Basketball is not known for its 'lifting culture', yet most athletes perform a warm-up before going on the court for a player development session. This an opportunity to do specific tissue loading based on athletic/injury profiles which serves to increase specific tissue capacity in a time efficient manner. The micro-dosed program can then be performed right after specific tissue loading or after the court workout. The actual training program should take from 15-20 minutes and focus on low volume, high intensity for the compound exercise(s) (Cuthbert et al., 2024). This helps provide an adequate stimulus without extraneous stress. The adequate stimulus means staving off the tissue and joint damage brought on through the multi-dimensional stressors of the NBA schedule.

1. **Addressing the Mechanical Governor:** If and when data from LPS monitoring shows a drop in metrics such as Braking Density between games, it is rarely an issue of true strength. Rather, it may be an issue of inhibition or reduction in neuromuscular capacity. For example, we know that as de-training occurs, which may inadvertently happen through the course of the season, muscles do not work as efficiently and cause inhibition. Similarly, fatigue caused in and between games may cause dysfunctional firing of muscles which increases injury risk (Hewett et al., 2005). Lastly, it is important to look at the ratios between accelerations and decelerations performed in games, as a decrease from

normative values may point to inhibition or neuromuscular fatigue. This is where the in-season strength and conditioning program comes into play. Through keeping a strength stimulus in between games, this help maintain proper firing patterns and tissue capacity.

2. **The Strength and Conditioning Program:** To explore this further, we can look at how stress accumulates on the body through games. Each game can be seen as a specific type of stress, filled with high-speed running (HSR), accelerations, and decelerations. While this stimulus is great for the sport of Basketball, it becomes too specific and repetitive of a stress on tissues. For example, HSR stresses the hamstrings as decelerations stress the quadriceps. While this is one way of increasing tissue capacity, there are other, more joint sparing ways of achieving this goal. By keeping a strength stimulus that emphasizes different movement patterns and compound ones such as squat variations, deadlift variations, etc. we get to promote strength while paradoxically avoiding additional stress on the body through movement variability (Bartlett et al., 2007).
3. **In-Season Strength and Conditioning:** Although the strength and conditioning culture in Basketball has historically been disparaging, this is partly because of incorrect assumptions. Modern strength and conditioning is carried out by emotionally intelligent, compassionate, and open-minded coaches (Nash & Collins, 2006). This translates into keeping the principles of strength and tissue capacity, while being able to listen and accept what types of movements athletes respond best to. In general, volume is low-moderate while intensity is high which allows for time efficient, yet substantial physical and physiological adaptations. In the stress of the season, this is one important aspect of keeping athletes active and healthy.

Conclusion

The Fatigue Paradox experienced throughout the NBA season cannot be solved by resting players alone. The thought that players should either not strength train during or do it sparingly through the season is outdated. Rather, players need to participate in a strength and conditioning to reduce DOMS, stay neuromuscularly fresh, and have a high level of tissue resilience. From the performance standpoint, the above points speak to this. However, there is a sports medicine

application piece to the in-season training sessions. Training sessions allow for specific tissue loading exercises to be programmed in, either as standalone exercises or accessory exercises. In this capacity, the performance and sports medicine staff are integrated and working together to increase local and global capacity. In-season lifting is already part of virtually every professional Basketball's club high performance plan, but it can be further optimized to increase player availability and longevity.

References

1. **Caparrós, T., et al. (2024).** Low External Workloads Are Related to Higher Injury Risk in Professional Male Basketball Games. *PubMed Central*.
2. **Cuthbert, M., et al. (2024).** The Braking Performance Framework: Practical Recommendations for Horizontal Deceleration. *International Journal of Strength and Conditioning*.
3. **Gabbett, T. J. (2023).** The Training-Injury Prevention Paradox: 10 Years Later. *British Journal of Sports Medicine*.
4. **Goitz, et al. (2023).** Neuromuscular Control and Proprioception in Elite Athletes: A Longitudinal Review. *Journal of Orthopaedic Research*.
5. **NBA/IQVIA Surveillance & Analytics. (2024).** Internal Report on Player Availability and Load Management.
6. **Huyghe, T., Alcaraz, P. E., Calleja-González, J., & Bird, S. P. (2022).** The underpinning factors of NBA game-play performance: A systematic review (2001–2020). *The Physician and Sportsmedicine*, 50(2), 94-122.
7. **Mandić, R., Jakovljević, S., Erčulj, F., & Štrumbelj, E. (2019).** Trends in NBA and Euroleague basketball: Analysis and comparison of statistical data from 2000 to 2017. *PloS one*, 14(10), e0223524.
8. **Bartolomei, S., Hoffman, J. R., Merni, F., & Stout, J. R. (2014).** A comparison of traditional and block periodized strength training programs in trained athletes. *The Journal of Strength & Conditioning Research*, 28(4), 990-997.
9. **Chiu, L. Z., & Barnes, J. L. (2003).** The fitness-fatigue model revisited: Implications for planning short-and long-term training. *Strength & Conditioning Journal*, 25(6), 42-51.
10. **Williams, S., West, S., Cross, M. J., & Stokes, K. A. (2017).** Better way to determine the acute: chronic workload ratio?. *British journal of sports medicine*, 51(3), 209-210.
11. **Vázquez-Guerrero, J., Suarez-Arrones, L., Gómez, D. C., & Rodas, G. (2018).** Comparing external total load, acceleration and deceleration outputs in elite basketball players across positions during match play. *Kinesiology*, 50(2), 228-234.
12. **Bishop, C., Jordan, M., Torres-Ronda, L., Loturco, I., Harry, J., Virgile, A., ... & Comfort, P. (2023).** Selecting metrics that matter: comparing the use of the countermovement jump for performance profiling, neuromuscular fatigue monitoring, and injury rehabilitation testing. *Strength & Conditioning Journal*, 45(5), 545-553.
13. **Cuthbert, M., Haff, G. G., McMahon, J. J., Evans, M., & Comfort, P. (2024).** Microdosing: a conceptual framework for use as programming strategy for resistance training in team sports. *Strength & Conditioning Journal*, 46(2), 180-201.
14. **Hewett, T. E., Zazulak, B. T., Myer, G. D., & Ford, K. R. (2005).** A review of electromyographic activation levels, timing differences, and increased anterior cruciate ligament injury incidence in female athletes. *British journal of sports medicine*, 39(6), 347-350.
15. **Bartlett, R., Wheat, J., & Robins, M. (2007).** Is movement variability important for sports biomechanists?. *Sports biomechanics*, 6(2), 224-243.
16. **Nash, C., & Collins, D. (2006).** Tacit knowledge in expert coaching: Science or art?. *Quest*, 58(4), 465-477.

Citation: Serrano B (2026); *The Fatigue Paradox: A Dual-Perspective Analysis of In-Season Neuromuscular Integrity and Load Management in the National Basketball Association*. *Adv Ortho and Sprts Med: AOASM-224*